



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Schwartz
Serial No. 09/815,978
Filed: March 22, 2001
For: *HYDRAZINE-BASED AND CARBONYL-
BASED BIFUNCTIONAL CROSSLINKING
REAGENTS*
Art Unit: Unassigned
Examiner: Unassigned

I hereby certify that this paper and the attached
papers are being deposited with the United States
Postal Service as first class mail in an envelope
addressed to:
Commissioner for Patents
Washington, D.C. 20231, on this date.

05/11/01
Date

Kelley Fischer
Kelly Fischer

PRELIMINARY AMENDMENT

Commissioner for Patents
Washington, D.C. 20231

Dear Sir:

Preliminary to examination of the above-captioned patent application,
please amend the application as follows:

IN THE SPECIFICATION:

Please amend the specification as follows (a marked-up copy of the
amended specification is attached to this Amendment):

**Please replace the paragraph on page 1, lines 10-17, with the following
paragraph.**

BACKGROUND OF THE INVENTION

Methods to crosslink biomolecules such as proteins, oligonucleotides and
carbohydrates to each other, to radioactive and non-radioactive metal chelates,
to drugs and to surfaces have allowed development of both in vitro and in vivo
diagnostic assays as well as in vivo therapies. A wide variety of methods have
been developed and reviewed (Greg T. Hermanson, Bioconjugate Techniques,
Academic Press).

U.S.S.N 09/815,978
Schwartz
PRELIMINARY AMENDMENT

Please replace the paragraph on page 11, lines 1-16, with the following paragraph.

As used herein, alkyl, alkenyl and alkynyl carbon chains, if not specified, contain from 1 to 20 carbons, preferably 1 to 16 carbons, and are straight or branched. Alkenyl carbon chains of from 2 to 20 carbons preferably contain 1 to 8 double bonds, and the alkenyl carbon chains of 1 to 16 carbons preferably contain 1 to 5 double bonds. Alkynyl carbon chains of from 2 to 20 carbons preferably contain 1 to 8 triple bonds, and the alkynyl carbon chains of 2 to 16 carbons preferably contain 1 to 5 triple bonds. Exemplary alkyl, alkenyl and alkynyl groups herein include, but are not limited to, methyl, ethyl, propyl, isopropyl, isobutyl, n-butyl, sec-butyl, tert-butyl, isopentyl, neopentyl, tert-pentyl and isohexyl. The alkyl, alkenyl and alkynyl groups, unless otherwise specified, can be optionally substituted, with one or more groups, preferably alkyl group substituents that can be the same or different. As used herein, lower alkyl, lower alkenyl, and lower alkynyl refer to carbon chains having less than about 6 carbons. As used herein, "alk(en)(yn)yl" refers to an alkyl group containing at least one double bond and at least one triple bond.

Please replace the paragraph beginning on page 11, line 26, through page 12, line 12, with the following paragraph.

As used herein, an "aryl group substituent" includes alkyl, cycloalkyl, cycloalkylalkyl, aryl, heteroaryl optionally substituted with 1 or more, preferably 1 to 3, substituents selected from halo, haloalkyl and alkyl, aralkyl, heteroaralkyl, alkenyl containing 1 to 2 double bonds, alkynyl containing 1 to 2 triple bonds, alk(en)(yn)yl groups, halo, pseudohalo, cyano, hydroxy, haloalkyl and polyhaloalkyl, preferably halo lower alkyl, especially trifluoromethyl, formyl, alkylcarbonyl, arylcarbonyl that is optionally substituted with 1 or more, preferably 1 to 3, substituents selected from halo, haloalkyl and alkyl, heteroarylcarbonyl, carboxy, alkoxycarbonyl, aryloxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylaminocarbonyl,

U.S.S.N 09/815,978
Schwartz
PRELIMINARY AMENDMENT

diarylaminocarbonyl, aralkylaminocarbonyl, alkoxy, aryloxy, perfluoroalkoxy, alkenyloxy, alkynyloxy, arylalkoxy, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, arylaminoalkyl, amino, alkylamino, dialkylamino, arylamino, alkylaryl amino, alkylcarbonylamino, arylcarbonylamino, azido, nitro, mercapto, alkylthio, arylthio, perfluoroalkylthio, thiocyano, isothiocyano, alkylsulfinyl, alkylsulfonyl, arylsulfinyl, arylsulfonyl, aminosulfonyl, alkylaminosulfonyl, dialkylaminosulfonyl and arylaminosulfonyl.

Please replace the paragraph beginning on page 12, line 30, through page 13, line 8, with the following paragraph.

As used herein, "heteroaryl" refers to a monocyclic or multicyclic ring system, preferably of about 5 to about 15 members where one or more, more preferably 1 to 3 of the atoms in the ring system is a heteroatom, that is, an element other than carbon, for example, nitrogen, oxygen and sulfur atoms. The heteroaryl can be optionally substituted with one or more, preferably 1 to 3, aryl group substituents. The heteroaryl group can be optionally fused to a benzene ring. Exemplary heteroaryl groups include, for example, furyl, imidazolyl, pyrrolidinyl, pyrimidinyl, tetrazolyl, thienyl, pyridyl, pyrrolyl, N-methylpyrrolyl, quinolinyl and isoquinolinyl, with pyridyl and quinolinyl being preferred.

Please replace the paragraph on page 24, lines 19-26, with the following paragraph.

In another embodiment, the bifunctional hydrazide reagents provided herein form acid cleavable hydrazones. These reagents are advantageous as they can be used to modify biomolecules or carriers such as polymers in a single step. These modified aliphatic hydrazide biomolecules or carriers can subsequently react with carbonyl containing biomolecules, drug or other therapeutic or diagnostic reagent to readily form a hydrazone that can be cleaved following exposure to mild aqueous acid conditions at pH < 5.

U.S.S.N 09/815,978
Schwartz
PRELIMINARY AMENDMENT

Please replace the paragraph on page 33, lines 10-24, with the following paragraph.

The development of both DNA-based and protein microarrays has led to a revolution in biotechnology. These microarrays are based on immobilization of tens to tens of thousand biomolecules on solid surfaces. Silica based surfaces such as glass slides and silica chips have been the surface of choice to prepare microarrays. The immobilization of biomolecules requires attachment of the biomolecules via covalent or non-covalent, i.e., electrostatic, interactions. Glass slides modified to incorporate amino or aldehyde groups are commercially available (www.arrayit.com, Telechem, Inc, Sunnyvale, CA and www.cel-1.com, Cel Associates, Houston, TX). Protocols to immobilize oligonucleotides or polynucleotides require the use of strong chemical conditions such as sodium borohydride or crosslinking conditions such as photolysis. These methods are inefficient and cause direct modification of the oligonucleotide leading to reduced affinity towards its complementary target.

Please replace the paragraphs on page 38, lines 8-15, with the following paragraphs.

Bifunctional carbazides or thiocarbazides may be prepared by treatment of a hydrazine with phosgene or thiophosgene, respectively, in the presence of base followed by isolation of the iso(thio)cyanate. Addition of hydrazine yields the desired carbazide or thiocarbazide respectively.

Bifunctional semicarbazides or thiosemicarbazides may be prepared by treatment of an amine with phosgene or thiophosgene, respectively, in the presence of base followed by isolation of the iso(thio)cyanate. Addition of hydrazine yields the desired semicarbazide or thiosemicarbazide respectively.

U.S.S.N 09/815,978
Schwartz
PRELIMINARY AMENDMENT

Please replace the paragraphs beginning on page 41, line 18, through page 42, line 4, with the following paragraphs.

To a solution of this residue in DMF is added t-butyl carbazate (1.0 equivalent; Aldrich Chemical Company, Milwaukee, WI) in DMF. The reaction mixture is stirred at room temperature for 2 hours. The solvent is removed under reduced pressure and the residue is partitioned between ethyl acetate and 5% aqueous citric acid. The organic phase is washed with brine, dried over magnesium sulfate, filtered and concentrated to give 4-(tert-butoxycarbonylthiosemicarbazidomethyl)cyclohexane carboxylic acid.

This compound (1.0 equivalent) is dissolved in DMF and N-hydroxysuccinimide (1.0 equivalent) is added followed by the dropwise addition of a solution of dicyclohexylcarbodiimide (1.0 equivalents) in DMF. The reaction mixture is stirred at room temperature for 4 hours. The dicyclohexylurea (DCU) precipitate byproduct is removed by filtration and the filtrate is concentrated to dryness. The residue is partitioned between ethyl acetate and 5% aqueous citric acid. The organic phase is washed with brine, dried over magnesium sulfate, filtered and concentrated to give succinimidyl 4-(tert-butoxycarbonylthiosemicarbazidomethyl)cyclohexane carboxylate.

Please replace the paragraph on page 42, lines 21-31, with the following paragraph.

This hydrazone (1.0 equivalent) was suspended in DMF and N-hydroxysuccinimide (NHS)(1.0 equivalent) was added and followed by the addition of a solution of DCC (1.0 equivalent) in DMF. The reaction mixture was stirred at room temperature for 16 hours. The heterogeneous reaction mixture was filtered and the filtrate was concentrated under reduced pressure. The residue was dissolved in a minimum amount of ethyl acetate and hexanes were added to turbidity. A pale yellow precipitate formed that was isolated by filtration to give the desired compound with an approximate yield of 33%. PMR

U.S.S.N 09/815,978
Schwartz
PRELIMINARY AMENDMENT

(DMSO-d₆) δ 1.99 (s, 3H), 2.00 (s, 3H), 3.32 (s, 4H), 7.17 (d, 1H), 8.12 (dd, 1H), 8.76 (d, 1H), 10.39 (s, 1H).

Please replace the paragraph on page 43, lines 12-18, with the following paragraph.

To a solution of the resulting compound (1 mmol) in THF is added maleic anhydride (1 mmol), the reaction mixture is stirred at room temperature and acetic anhydride (1 mmol) and triethylamine (1 mmol) are added. Following stirring at room temperature for 16 hours, the solvent is removed under reduced pressure, and the residue is chromatographed on silica gel using ethyl acetate as eluant. The fractions containing product are pooled and concentrated.

Please replace the paragraph beginning on page 43, line 24, through page 44, line 3, with the following paragraph.

EXAMPLE 4

Preparation of Carbocyclic Succinimidyl Thiosemicarbazide Hydrochloride

To a suspension of proline (1 mmol) in THF is added triethylamine (2.5 mmol) followed by the dropwise addition of a solution of thiophosgene (1.1 mmol). The reaction mixture is stirred at ambient temperature for 4 hours followed by cooling the reaction mixture to 0 °C and the dropwise addition of a solution of t-butyl carbazate (1.1 mL). The reaction mixture is stirred at 0 °C for 1 hour and at room temperature for 2 hours. The solvent is removed under reduced pressure, and the residue is chromatographed on silica gel using methylene chloride/methanol (9/1) as eluant. The fractions containing product are pooled and concentrated.

Please replace the paragraphs beginning on page 46, line 2, through page 47, line 7, with the following paragraphs.

EXAMPLE 7

Preparation of 96 Well Plates to Incorporate Aromatic Aldehyde Moieties

U.S.S.N 09/815,978
Schwartz
PRELIMINARY AMENDMENT

Amino-modified 96 well plates (Costar or Corning) are modified with succinimidyl 4-formylbenzoate (SFB) as follows. A fresh solution of SFB (1.25 mL of 10 mg/mL) in DMSO is prepared. This solution is diluted into phosphate buffered saline (PBS) (0.1 M phosphate, 0.15 M NaCl, pH 7.4: 100 mL). To each well is added 200 μ L of the SFB/PBS solution and the wells are incubated at room temperature for 4 hours. The wells are washed three times with PBS/0.5% Tween®. The wells are dried and are ready for protein conjugation.

EXAMPLE 8

Preparation of 96 Well Plates to Incorporate Aromatic Hydrazine Moieties

Amino-modified 96 well plates (Costar or Corning) are modified with succinimidyl acetone nicotinic acid hydrazone (SANH) as follows. A fresh solution of SANH (1.25 mL of 10 mg/mL) in DMSO is prepared. This solution is diluted into PBS (0.1 M phosphate, 0.15 M NaCl, pH 7.4: 100 mL). To each well was added 200 μ L of the SANH/PBS solution, and the wells were incubated at room temperature for 4 hours. The wells are washed with water and then treated with 0.1 M acetate, pH 4.7 (200 μ L) for 2 hours. The wells were washed three times with PBS/0.5% Tween®. The wells were dried and are ready for conjugation to molecules possessing carbonyl moieties.

EXAMPLE 9

General Procedure for the Modification of a Protein with a Succinimidyl Carbazidyl Modification Reagent

A 5 mg/mL solution of bovine serum albumin in PBS (100 mM phosphate, 150 mM NaCl, pH 7.4 and 2 mM EDTA) (200 μ L; 1 mg protein) is prepared. A solution of succinimidyl 4-semicarbazidylbenzoate hydrochloride (SSCH; 3.5 mg) in DMF (100 μ L) is prepared. To the protein solution is added the SSCH/DMF solution (30 equivalents). The reaction mixture is incubated at room temperature for 4 hours. The modified protein is isolated by placing the reaction mixture in a 30K ultra-free centrifugation device and washing three times with conjugation buffer (3 X 400 μ L). The purified protein is quantitated for protein

U.S.S.N 09/815,978
Schwartz
PRELIMINARY AMENDMENT

concentration (BCA assay) and for hydrazine modification level by addition of 0.2 mM 2-p-nitrobenzaldehyde in PBS pH 7.4 and measuring the absorbance at 380nm (extinction coefficient 22,600).

Please replace the paragraphs beginning on page 47, line 25, through page 49, line 30, with the following paragraphs.

EXAMPLE 11

General procedure for preparation of a hydrazine-modified protein

A 5 mg/mL solution of ovalbumin in PBS (100 mM phosphate, 150 mM NaCl, pH 7.4) and 2 mM EDTA (200 μ L; 1 mg protein) was prepared. A solution of succinimidyl 6-hydrazinonicotinate acetone hydrazone (SANH)(EXAMPLE 2)(2 mg) in DMF (50 μ L) is prepared. To the protein solution was added the SANH/DMF solution (15 eq.). The reaction mixture was incubated at room temperature for 4 hours. The modified protein was isolated and buffer exchanged by placing the reaction mixture in a 30K ultra-free centrifugation device and washing three times with 0.1 M MES, 0.9% NaCl, pH 4.7 (3 X 400 μ L). The purified protein was quantified for protein concentration (BCA assay; Pierce Chemical Co., Rockford, IL) and for hydrazine modification level by incubation of an aliquot of protein in a 0.5 mM 4-nitrobenzaldehyde in 0.1 M MES, 0.9% NaCl, pH 4.7 and measuring the absorbance at 360 nm (molar extinction coefficient 22,000).

EXAMPLE 12

General Procedure for the Preparation of a Conjugate by Reaction of a Hydrazine Modified Protein with a Carbonyl Modified Protein

Aldehyde-modified IgG (EXAMPLE 10) in MES (1 mg; 0.200 μ L of a 2.5 mg/mL solution) was added to a solution of hydrazine-modified ovalbumin (EXAMPLE 11, 1 mL; 0.200 μ L of a 5 mg/mL solution), and the reaction mixture was incubated at room temperature for 4 hours. The reaction mixture was analyzed by PAGE gel (coomassie blue development) that demonstrated presence of a high molecular weight product and <5% unreacted aldehyde-

U.S.S.N 09/815,978
Schwartz
PRELIMINARY AMENDMENT

modified IgG and <10% unreacted hydrazine-modified ovalbumin. The level of conjugation is quantified by measuring the absorbance at 360 nm.

EXAMPLE 13

Preparation of a thiosemicarbazide modified protein

A 5 mg/mL solution of ovalbumin in PBS (100 mM phosphate, 150 mM NaCl, pH 7.4) and 2 mM EDTA (200 μ L; 1 mg protein) was prepared. A solution of succinimidyl 4-thiosemicarbazidylbenzoate hydrochloride (STBH)(2 mg) in DMF (50 μ L) is prepared. To the protein solution was added the STBH/DMF solution (15 eq.). The reaction mixture was incubated at room temperature for 4 hours. The modified protein was isolated and buffer exchanged by placing the reaction mixture in a 30K ultra-free centrifugation device and washing three times with 0.1 M MES, 0.9% NaCl, pH 4.7 (3 X 400 μ L). The purified protein was quantified for protein concentration (BCA assay; Pierce Chemical Co., Rockford, IL).

EXAMPLE 14

Conjugation of a thiosemicarbazide-modified protein to an aldehyde modified protein

The thiosemicarbazide protein prepared in EXAMPLE 13 was reacted with aldehyde-modified protein in an identical manner as described for the hydrazine-modified protein in EXAMPLE 12. Analysis by PAGE gel demonstrated similar efficiency as conjugation as observed in EXAMPLE 12.

EXAMPLE 15

Preparation of a hydrazide modified protein

A 5 mg/mL solution of ovalbumin in PBS (100 mM phosphate, 150 mM NaCl, pH 7.4) and 2 mM EDTA (200 μ L; 1 mg protein) was prepared. A solution of succinimidyl 4-hydrazidoterephthalate hydrochloride (SHTH)(2 mg) in DMF (50 μ L) is prepared. To the protein solution was added the SHTH/DMF solution (15 eq.). The reaction mixture was incubated at room temperature for 4 hours. The modified protein was isolated and buffer exchanged by placing the reaction mixture in a 30K ultra-free centrifugation device and washing three times with

U.S.S.N 09/815,978
Schwartz
PRELIMINARY AMENDMENT

0.1 M MES, 0.9% NaCl, pH 4.7 (3 X 400 μ L). The purified protein was quantified for protein concentration (BCA assay; Pierce Chemical Co., Rockford, IL).

EXAMPLE 16

Conjugation of a hydrazide-modified protein to an aldehyde modified protein

The hydrazide-modified protein prepared in EXAMPLE 15 was reacted with aldehyde-modified protein in an identical manner as described for the hydrazine-modified protein in EXAMPLE 12. Analysis by PAGE gel demonstrated similar efficiency as conjugation as observed in EXAMPLE 12.

Please replace the paragraph on page 51, lines 14-22, with the following paragraph.

EXAMPLE 21

Immobilization of Oxidized Horseradish peroxidase to Hydrazine-Modified Plates

Periodate-oxidized horseradish peroxidase (Pierce Chemical Co., Rockford, IL) is diluted to the desired concentration and added to 96-3456 well plates that had been modified to possess hydrazino groups as described above in EXAMPLE 9. The antibody solution is allowed to incubate for 2-18 hours followed by removal of the solution and washing with 0.5% Tween[®] solution (twice) and buffer (twice).

Please replace the paragraph beginning on page 53, line 18, through page 54, line 8, with the following paragraph.

EXAMPLE 26

Preparation of a bifunctional aromatic oxyammonium hydrochloride, -O-oxyammoniumnicotinate hydrochloride (SONH)

As shown in FIGURE 9, to a solution of 6-chloronicotinic acid (1 equivalent) in 80% aqueous ethanol is added hydroxylamine (500 equivalents) and the solution is refluxed for 16 hours. The reaction mixture is concentrated to dryness and dissolved in water. The solution cooled in an ice bath and

U.S.S.N 09/815,978
Schwartz
PRELIMINARY AMENDMENT

acidified with concentrated hydrochloric acid until a precipitate forms, pH approximately 5.0. The solids are isolated, redissolved in water and the pH of the solution raised to 7.5 with base. Dioxane (1 volume) is added to the solution followed by the dropwise addition of di-t-butyl dicarbonate (5 equivalents; Aldrich Chemical Co.). The reaction mixture is stirred at room temperature for 4 hours and the dioxane removed on the rotavap. The residue is chromatographed on silica to isolate the desired BOC acid. The acid (1 equiv) is dissolved in DMF and treated with NHS (1 equiv) followed by the dropwise addition of DCC (1 equiv) in DMF. The reaction mixture is stirred at room temperature for 4 hours and the solids removed by filtration and the filtrate concentrated to dryness and resuspended in ethyl acetate. Further precipitate is removed by filtration and the filtrate concentrated to dryness and the desired BOC succinimidyl ester is isolated by silica gel chromatography.

Please replace the paragraphs beginning on page 54, line 18, through page 55, line 30, with the following paragraphs.

EXAMPLE 27

Preparation of an oxyamino-modified protein

A 5 mg/mL solution of ovalbumin in PBS (100 mM phosphate, 150 mM NaCl, pH 7.4) and 2 mM EDTA (200 μ L; 1 mg protein) is prepared. A solution of succinimidyl aminooxyacetate hydrochloride (SAAH)(2 mg) in DMF (50 μ L) is prepared. To the protein solution is added the SAAH/DMF solution (15 eq.). The reaction mixture is incubated at room temperature for 4 hours. The modified protein is isolated and buffer exchanged by placing the reaction mixture in a 30K ultra-free centrifugation device and washing three times with 0.1 M MES, 0.9% NaCl, pH 4.7 (3 X 400 μ L). The purified protein is quantified for protein concentration (BCA assay; Pierce Chemical Co., Rockford, IL).

EXAMPLE 28

Preparation of a hydrazinonicotinamide modified polymer

A solution of poly-L-lysine (10 mg; Sigma Chemicals, St. Louis, MO; cat. #P-7890) was dissolved in conjugation buffer, 0.1 M phosphate, 0.15 M NaCl, pH 7.4 (1 mL). A solution of succinimidyl 6-hydrazinonicotinate acetone hydrazone (SANH; 1.3 mg) was dissolved in DMSO (13 μ L). To two poly-L-lysine aliquots (200 μ L) were added the SANH/DMSO solution (2.85 μ L (10 equivalents) and 5.7 μ L (20 equivalents)). The reaction mixtures were vortexed and incubated at room temperature for 2 h. The modified polymer was isolated by gel filtration on a NAP-25 column (Pharmacia) pre-equilibrated with 0.1 M MES, 0.9% NaCl, pH 4.7 buffer. Fractions (1 mL) were collected and analyzed by UV (A260). Fractions containing UV active product were combined to yield the desired product. The product was analyzed colorimetrically for hydrazine content by dissolving an aliquot (2 μ L) in a 0.5 mM solution of p-nitrobenzaldehyde (98 μ L) and incubating at 37 °C for 1 h followed by taking A390 readings (extinction coefficient 22000). The HyNic:poly-L-lysine polymer was used directly in the conjugation step. The amine/hydrazine content was determined using the TNBSA assay (trinitrobenzenesulfonic acid; Pierce Chemical, Inc., Rockville, IL).

EXAMPLE 29

Preparation of an aliphatic hydrazine modified protein and conjugation to periodate oxidized bacterial polysaccharide

A solution of a bacterial polysaccharide that possesses unsaturation in its lipids (from ATCC; 10 mg/mL) in water is treated with 10 mM sodium periodate (1/10 volume to make the solution 1 mM in periodate) and incubated at room temperature for 30 min. The reaction mixture is passed through a sephadex G-25 column pre-equilibrated with water to remove small molecule impurities. The polysaccharide combining fractions are combined and concentrated to 5 mg/mL.

U.S.S.N 09/815,978
Schwartz
PRELIMINARY AMENDMENT

IN THE CLAIMS:

Please replace claims 12, 23, 24, 35, 37, and 49-53 with the following amended claims (a marked up copy of the amended claims is attached to this Amendment):


12. (Amended) The compound of claim 10, wherein R is a divalent aliphatic group.

23. (Amended) The compound of claim 1, wherein B is an amino reactive moiety selected from succinimidyl ester, hydroxybenzotriazolyl ester, or pentafluorophenol ester.

24. (Amended) The compound of claim 1, wherein B is a thiol reactive moiety selected from maleimido, α -bromoacetyl, α -bromoacetamido or pyridyldisulfide.

35. (Amended) A method of crosslinking a natural or synthetic biological molecule, comprising:

(i) preparing a conjugate of formula Va:



or a derivative thereof, wherein:

A is NH(C=O), NH(C=S), NH(C=NH), NHNH(C=O), NHNH(C=S), NHNH(C=NH) or a direct bond;

B is a natural or synthetic biological molecule;

D is a carbon or nitrogen atom;

E is a carbon or nitrogen atom;

R¹ is hydrogen or a saturated straight chain of 1 to 12 carbon atoms; and

R² is hydrogen or a saturated straight chain of 1 to 12 carbon atoms; and


(ii) applying the conjugate to a surface wherein the surface has at least one carbonyl moiety for a time and under conditions such that the

U.S.S.N 09/815,978
Schwartz
PRELIMINARY AMENDMENT

hydrazine moiety of the conjugate reacts with the carbonyl moiety of the surface forming a hydrazone bond to the surface.

37. (Amended) A method of crosslinking a natural or synthetic biological molecule, comprising:

(i) preparing a conjugate of formula IVa:



or a derivative thereof, wherein:

A is $\text{NH}(\text{C}=\text{O})$, $\text{NH}(\text{C}=\text{S})$, $\text{NH}(\text{C}=\text{NH})$, $\text{NHNH}(\text{C}=\text{O})$, $\text{NHNH}(\text{C}=\text{S})$, $\text{NHNH}(\text{C}=\text{NH})$ or a direct bond;

B is a natural or synthetic biological molecule;

D is a carbon or nitrogen atom;

E is a carbon or nitrogen atom; and

X is a negative counter ion, oxygen, sulfur or $-\text{NH}$; and

(ii) mixing the conjugate to a natural or synthetic biological molecule, wherein the molecule has at least one carbonyl moiety, for a time and under conditions such that the hydrazine moiety of the conjugate reacts with the carbonyl moiety of the molecule forming a hydrazone bond to the molecule.

49. (Amended) The compound of claim 5, wherein B is an amino reactive moiety selected from succinimidyl ester, hydroxybenzotriazolyl ester, or pentafluorophenol ester.

50. (Amended) The compound of claim 8, wherein B is an amino reactive moiety selected from succinimidyl ester, hydroxybenzotriazolyl ester, or pentafluorophenol ester.

51. (Amended) The compound of claim 10, wherein B is an amino reactive moiety selected from succinimidyl ester, hydroxybenzotriazolyl ester, or pentafluorophenol ester.

U.S.S.N 09/815,978
Schwartz
PRELIMINARY AMENDMENT

52. (Amended) The compound of claim 5, wherein B is a thiol reactive moiety selected from maleimido, α -bromoacetyl, α -bromoacetamido or pyridyldisulfide.

53. (Amended) The compound of claim 10, wherein B is a thiol reactive moiety selected from maleimido, α -bromoacetyl, α -bromoacetamido or pyridyldisulfide.

REMARKS

Any fees that may be due in connection with this application throughout its pendency may be charged to Deposit Account No. 50-1213.

The specification is amended to correct obvious typographical and spelling errors and to produce grammatical clarity. The specification is also amended to correct the name of a chemical compound in the paragraph on page 38, lines 12-15. The basis for this amendment is found in the specification, in particular page 38, line 12, which describes the preparation of semicarbazides and thiosemicarbazides. The specification is also amended to correct an error in the solvent description found in the paragraphs on page 41, line 18 to page 42, line 4. This amendment finds basis in Figure 3 which shows that the reactions are done in DMF as the solvent. The specification is also amended to correct the name of a buffer in the paragraph on page 46, lines 12-23. The basis for this amendment is found in the specification, in particular page 46, lines 16-17, which describes the preparation of a "SANH/PBS" solution.

The amendments to claims 12, 23, and 49-51 correct obvious typographical and spelling errors. The amendments to claims 24, 52, and 53 correct typographical errors and find basis in the specification, in particular page 4, line 28, which describes the α -bromoacetamido group as a thiol reactive moiety. Claims 32 and 35 are identical. The amendment to claim 35 corrects this redundancy and finds basis in the specification, in particular, page 20, lines 11-15, which describes methods that provide conjugates. The amendment to claim 37 corrects a chemical structure and finds basis in the specification, in

U.S.S.N 09/815,978
Schwartz
PRELIMINARY AMENDMENT

particular page 26, line 5, which shows the chemical structure of formula IVa.
No new matter has been added.


Included as an attachment is a marked-up version of the specification paragraphs that are being amended, per 37 CFR §1.121.

* * *

Entry of this amendment and examination of the application are respectfully requested.

Respectfully submitted,
HELLER EHRMAN WHITE & McAULIFFE LLP

By:


Stephanie L. Seidman
Registration No. 33,779

Attorney Docket No. 37154-753
Address all correspondence to:
Stephanie L. Seidman, Esq.
HELLER EHRMAN WHITE & McAULIFFE
4350 La Jolla Village Drive, 6th Floor
San Diego, California 92122-1246
Telephone: 858 450-8400
Facsimile: 858 587-5360
email:sseidman@HEWM.com



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Schwartz
Serial No. 09/815,978
Filed: March 22, 2001
For: *HYDRAZINE-BASED AND CARBONYL-
BASED BIFUNCTIONAL CROSSLINKING
REAGENTS*
Art Unit: Unassigned
Examiner: Unassigned

ATTACHMENT TO THE PRELIMINARY AMENDMENT
MARKED UP PARAGRAPHS AND CLAIMS (37 CFR §1.121)

IN THE SPECIFICATION:

Please amend the specification as follows:

Please amend the paragraph on page 1, lines 10-17, as follows:

BACKGROUND OF THE INVENTION

Methods to crosslink biomolecules such as proteins, oligonucleotides and carbohydrates to each other, to radioactive and non-radioactive metal chelates, to drugs and to surfaces [has] have allowed development of both in vitro and in vivo diagnostic assays as well as in vivo therapies. A wide variety of methods have been developed and reviewed (Greg T. Hermanson, Bioconjugate Techniques, Academic Press).

Please amend the paragraph on page 11, lines 1-16, as follows:

As used herein, alkyl, alkenyl and alkynyl carbon chains, if not specified, contain from 1 to 20 carbons, preferably 1 to 16 carbons, and are straight or branched. Alkenyl carbon chains of from 2 to 20 carbons preferably contain 1 to 8 double bonds, and the alkenyl carbon chains of 1 to 16 carbons preferably contain 1 to 5 double bonds. Alkynyl carbon chains of from 2 to 20 carbons preferably contain 1 to 8 triple bonds, and the alkynyl carbon chains of 2 to 16

PRELIMINARY AMENDMENT ATTACHMENT

carbons preferably contain 1 to 5 triple bonds. Exemplary alkyl, alkenyl and alkynyl groups herein include, but are not limited to, methyl, ethyl, propyl, isopropyl, isobutyl, n-butyl, sec-butyl, tert-butyl, isopentyl, neopentyl, tert-pen[y]tyl and isohexyl. The alkyl, alkenyl and alkynyl groups, unless otherwise specified, can be optionally substituted, with one or more groups, preferably alkyl group substituents that can be the same or different. As used herein, lower alkyl, lower alkenyl, and lower alkynyl refer to carbon chains having less than about 6 carbons. As used herein, "alk(en)(yn)yl" refers to an alkyl group containing at least one double bond and at least one triple bond.

Please amend the paragraph beginning on page 11, line 26, through page 12, line 12, as follows:

As used herein, an "aryl group substituent" includes alkyl, cycloalkyl, cycloalkylalkyl, aryl, heteroaryl optionally substituted with 1 or more, preferably 1 to 3, substituents selected from halo, halo[]alkyl and alkyl, aralkyl, heteroaralkyl, alkenyl containing 1 to 2 double bonds, alkynyl containing 1 to 2 triple bonds, alk(en)(yn)yl groups, halo, pseudohalo, cyano, hydroxy, haloalkyl and polyhaloalkyl, preferably halo lower alkyl, especially trifluoromethyl, formyl, alkylcarbonyl, arylcarbonyl that is optionally substituted with 1 or more, preferably 1 to 3, substituents selected from halo, halo[]alkyl and alkyl, heteroarylcarbonyl, carboxy, alkoxycarbonyl, aryloxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylaminocarbonyl, diarylaminocarbonyl, aralkylaminocarbonyl, alkoxy, aryloxy, perfluoroalkoxy, alkenyloxy, alkynyloxy, arylalkoxy, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, arylaminoalkyl, amino, alkylamino, dialkylamino, arylamino, alkylaryl amino, alkylcarbonylamino, arylcarbonylamino, azido, nitro, mercapto, alkylthio, arylthio, perfluoroalkylthio, thiocyno, isothiocyano, alkylsulfinyl, alkylsulfonyl, arylsulfinyl, arylsulfonyl, aminosulfonyl, alkylaminosulfonyl, dialkylaminosulfonyl and arylaminosulfonyl.

U.S.S.N 09/815,978
Schwartz
PRELIMINARY AMENDMENT ATTACHMENT

Please amend the paragraph beginning on page 12, line 30, through page 13, line 8, as follows:

As used herein, "heteroaryl" refers to a monocyclic or multicyclic ring system, preferably of about 5 to about 15 members where one or more, more preferably 1 to 3 of the atoms in the ring system is a heteroatom, that is, an element other than carbon, for example, nitrogen, oxygen and sulfur atoms. The heteroaryl can be optionally substituted with one or more, preferably 1 to 3, aryl group substituents. The heteroaryl group can be optionally fused to a benzene ring. Exemplary heteroaryl groups include, for example, furyl, imidaz[in]olyl, pyrrolidinyl, pyrimidinyl, tetrazolyl, thienyl, pyridyl, pyrrolyl, N-methylpyrrolyl, quinolinyl and isoquinolinyl, with pyridyl and quinolinyl being preferred.

Please amend the paragraph on page 24, lines 19-26, as follows:

In another embodiment, the bifunctional hydrazide reagents provided herein form acid cleavable hydrazones. These reagents are advantageous as they can be used to modify biomolecules or carriers such as polymers in a single step. These modified aliphatic hydrazide biomolecules or carriers can subsequently react[ed] with carbonyl containing biomolecules, drug or other therapeutic or diagnostic reagent to readily form a hydrazone that can be cleaved following exposure to mild aqueous acid conditions at pH < 5.

Please amend the paragraph on page 33, lines 10-24, as follows:

The development of both DNA-based and protein microarrays has led to a revolution in biotechnology. These microarrays are based on immobilization of tens to tens of thousand biomolecules on solid surfaces. Silica based surfaces such as glass slides and silica chips have been the surface of choice to prepare microarrays. The immobilization of biomolecules requires attachment of the biomolecules via covalent or non-covalent, i.e., electrostatic, interactions. Glass slides modified to incorporate amino or aldehyde groups are commercially

available (www.arrayit.com, Telechem, Inc, [someplace] Sunnyvale, CA and www.cel-1.com, Cel Associates, Houston, TX). Protocols to immobilize oligonucleotides or polynucleotides require the use of strong chemical conditions such as sodium borohydride or crosslinking conditions such as photolysis. These methods are inefficient and cause direct modification of the oligonucleotide leading to reduced affinity towards its complementary target.

Please amend the paragraphs on page 38, lines 8-15, as follows:

Bifunctional carbazides or thiocarbazides may be prepared by treatment of a hydrazine with phosgene or thiophosgene, respectively, in the presence of base followed by isolation of the iso(thio)cyanate. Addition of hydrazine yields the desired carbazide or thiocarbazide respectively.

Bifunctional semicarbazides or thiosemicarbazides may be prepared by treatment of an amine with phosgene or thiophosgene, respectively, in the presence of base followed by isolation of the iso(thio)cyanate. Addition of hydrazine yields the desired semicarbazide or thiosemicarbazide respectively.

Please amend the paragraphs beginning on page 41, line 18, through page 42, line 4, as follows:

To a solution of this residue in [THF] DMF is added t-butyl carbazate (1.0 equivalent; Aldrich Chemical Company, Milwaukee, WI) in [THF] DMF. The reaction mixture is stirred at room temperature for 2 hours. The solvent is removed under reduced pressure and the residue is partitioned between ethyl acetate and 5% aqueous citric acid. The organic phase is washed with brine, dried over magnesium sulfate, filtered and concentrated to give 4-(tert-butoxycarbonylthiosemicarbazidomethyl)cyclohexane carboxylic acid.

This compound (1.0 equivalent) is dissolved in [THF] DMF and N-hydroxysuccinimide (1.0 equivalent) is added followed by the dropwise addition of a solution of dicyclohexylcarbodiimide (1.0 equivalents) in [THF] DMF. The reaction mixture is stirred at room temperature for 4 hours. The

dicyclohexylurea (DCU) precipitate byproduct is removed by filtration and the filtrate is concentrated to dryness. The residue is partitioned between ethyl acetate and 5% aqueous citric acid. The organic phase is washed with brine, dried over magnesium sulfate, filtered and concentrated to give succinimidyl 4-(tert-butoxycarbonylthiosemicarbazidomethyl)cyclohexane carboxylate.

Please amend the paragraph on page 42, lines 21-31, as follows:

This hydrazone (1.0 equivalent) was suspended in DMF and N-hydroxysuccinimide (NHS)(1.0 equivalent) was added and followed by the addition of a solution of DCC (1.0 equivalent) in DMF[was added]. The reaction mixture was stirred at room temperature for 16 hours. The heterogeneous reaction mixture was filtered and the filtrate was concentrated under reduced pressure. The residue was dissolved in a minimum amount of ethyl acetate and hexanes were added to turbidity. A pale yellow precipitate formed that was isolated by filtration to give the desired compound with an approximate yield of 33%. PMR (DMSO-d₆) δ 1.99 (s, 3H), 2.00 (s, 3H), 3.32 (s, 4H), 7.17 (d, 1H), 8.12 (dd, 1H), 8.76 (d, 1H), 10.39 (s, 1H).

Please amend the paragraph on page 43, lines 12-18, as follows:

To a solution of the resulting compound (1 mmol) in THF is added maleic anhydride (1 mmol), the reaction mixture is stirred at room temperature and acetic anhydride (1 mmol) and triethylamine (1 mmol) [is] are added. Following stirring at room temperature for 16 hours, the solvent is removed under reduced pressure, and the residue is chromatographed on silica gel using ethyl acetate as eluant. The fractions containing product are pooled and concentrated.

Please amend the paragraph beginning on page 43, line 24, through page 44, line 3, as follows:

EXAMPLE 4

Preparation of Carbocyclic Succinimidyl Thiosemicarbazide Hydrochloride

To a suspension of proline (1 mmol) in THF is added triethylamine (2.5 mmol) followed by the dropwise addition of a solution of thiophosgene (1.1 mmol). The reaction mixture is stirred at ambient temperature for 4 hours followed by cooling the reaction mixture to 0 °C and the dropwise addition of a solution of t-butyl carbazate (1.1 mL). The reaction mixture is stirred at 0 °C for 1 hour and at room temperature for 2 hours. The solvent is removed under reduced pressure, and the residue is chromatographed on silica gel using methylene chloride/meth[ona]anol (9/1) as eluant. The fractions containing product are pooled and concentrated.

Please amend the paragraphs beginning on page 46, line 2, through page 47, line 7, as follows:

EXAMPLE 7

Preparation of 96 Well Plates to Incorporate Aromatic Aldehyde Moieties

Amino-modified 96 well plates (Costar or Corning) are modified with succinimidyl 4-formylbenzoate (SFB) as follows. A fresh solution of SFB (1.25 mL of 10 mg/mL) in DMSO is prepared. This solution is diluted into phosphate buffered saline (PBS)(0.1 M phosphate, 0.15 M NaCl, pH 7.4: 100 mL). To each well is added 200 μ L of the SFB/PBS solution and the wells are incubated at room temperature for 4 hours. The wells are washed three times with PBS/0.5% [t]Tween®. The wells are dried and are ready for protein conjugation.

EXAMPLE 8

Preparation of 96 Well Plates to Incorporate Aromatic Hydrazine Moieties

Amino-modified 96 well plates (Costar or Corning) are modified with succinimidyl acetone nicotinic acid hydrazone (SANH) as follows. A fresh solution of SANH (1.25 mL of 10 mg/mL) in DMSO is prepared. This solution is diluted into PBS (0.1 M phosphate, 0.15 M NaCl, pH 7.4: 100 mL). To each well was added 200 μ L of the S[FB]ANH/PBS solution, and the wells were incubated at room temperature for 4 hours. The wells are washed with water and then

treated with 0.1 M acetate, pH 4.7 (200 μ L) for 2 hours. The wells were washed three times with PBS/0.5% Tween[®]. The wells were dried and are ready for conjugation to molecules possessing carbonyl moieties.

EXAMPLE 9

General Procedure for the Modification of a Protein with a Succinimidyl Carbazidyl Modification Reagent

A 5 mg/mL solution of bovine serum albumin in PBS (100 mM phosphate, 150 mM NaCl, pH 7.4 and 2 mM EDTA) (200 μ L; 1 mg protein) is prepared. A solution of succinimidyl 4-semicarbazidylbenzoate hydrochloride (SSCH; 3.5 mg) in DMF (100 μ L) is prepared. To the protein solution is added the SSCH/DMF solution (30 equivalents). The reaction mixture is incubated at room temperature for 4 hours. The modified protein is isolated by placing the reaction mixture in a 30K ultra-free centrifugation device and washing three times with conjugation buffer (3 X 400 μ L). The purified protein is quantitated for protein concentration (BCA assay) and for hydrazine modification level by addition of 0.2 [M]mM 2-p-nitrobenzaldehyde in PBS pH 7.4 and measuring the absorbance at 380nm (extinction coefficient 22,600).

Please amend the paragraphs beginning on page 47, line 25, through page 49, line 30, as follows:

EXAMPLE 11

General procedure for preparation of a hydrazine-modified protein

A 5 mg/mL solution of ovalbumin in PBS (100 mM phosphate, 150 mM NaCl, pH 7.4) and 2 mM EDTA (200 μ L; 1 mg protein) was prepared. A solution of succinimidyl 6-hydrazinonicotinate acetone hydrazone (SANH)(EXAMPLE 2)(2 mg) in DMF (50 μ L) is prepared. To the protein solution was added the SANH/DMF solution (15 eq.). The reaction mixture was incubated at room temperature for 4 hours. The modified protein was isolated and buffer exchanged by placing the reaction mixture in a 30K ultra-free centrifugation

device and washing three times with 0.1 M MES, 0.9% NaCl, pH 4.7 (3 X 400 μ L). The purified protein was quantified for protein concentration (BCA assay; Pierce Chemical Co., Rockford, IL) and for hydrazine modification level by incubation of an aliquot of protein in a 0.5 mM 4-nitrobenzaldehyde in 0.1 M MES, 0.9% NaCl, pH 4.7 and measuring the absorbance at 360 nm (molar extinction coefficient 22,000).

EXAMPLE 12

General Procedure for the Preparation of a Conjugate by Reaction of a Hydrazine Modified Protein with a Carbonyl Modified Protein

Aldehyde-modified IgG (EXAMPLE 10) in MES (1 mg; 0.200 μ L of a 2.5 mg/mL solution) was added to a solution of hydrazine-modified ovalbumin (EXAMPLE 11, 1 mL; 0.200 μ L of a 5 mg/mL solution), and the reaction mixture was incubated at room temperature for 4 hours. The reaction mixture was analyzed by PAGE gel (coomassie blue development) that demonstrated presence of a high molecular weight product and <5% unreacted aldehyde-modified IgG and <10% unreacted hydrazine-modified ovalbumin. The level of conjugation is quantified by measuring the absorbance at 360 nm.

EXAMPLE 13

Preparation of a thiosemicarbazide modified protein

A 5 mg/mL solution of ovalbumin in PBS (100 mM phosphate, 150 mM NaCl, pH 7.4) and 2 mM EDTA (200 μ L; 1 mg protein) was prepared. A solution of succinimidyl 4-thiosemicarbazidylbenzoate hydrochloride (STBH)(2 mg) in DMF (50 μ L) is prepared. To the protein solution was added the STBH/DMF solution (15 eq.). The reaction mixture was incubated at room temperature for 4 hours. The modified protein was isolated and buffer exchanged by placing the reaction mixture in a 30K ultra-free centrifugation device and washing three times with 0.1 M MES, 0.9% NaCl, pH 4.7 (3 X 400 μ L). The purified protein was quantified for protein concentration (BCA assay; Pierce Chemical Co., Rockford, IL).

EXAMPLE 14

Conjugation of a thiosemicarbazide-modified protein to an aldehyde modified protein

The thiosemicarbazide protein prepared in EXAMPLE 13 was reacted with aldehyde-modified protein in an identical manner as described for the hydrazine-modified protein in EXAMPLE 12. Analysis by PAGE gel demonstrated similar efficiency as conjugation as observed in EXAMPLE 12.

EXAMPLE 15

Preparation of a hydrazide modified protein

A 5 mg/mL solution of ovalbumin in PBS (100 mM phosphate, 150 mM NaCl, pH 7.4) and 2 mM EDTA (200 μ L; 1 mg protein) was prepared. A solution of succinimidyl 4-hydrazidoterephthalate hydrochloride (SHTH)(2 mg) in DMF (50 μ L) is prepared. To the protein solution was added the SHTH/DMF solution (15 eq.). The reaction mixture was incubated at room temperature for 4 hours. The modified protein was isolated and buffer exchanged by placing the reaction mixture in a 30K ultra-free centrifugation device and washing three times with 0.1 M MES, 0.9% NaCl, pH 4.7 (3 X 400 μ L). The purified protein was quantified for protein concentration (BCA assay; Pierce Chemical Co., Rockford, IL).

EXAMPLE 16

Conjugation of a hydrazide-modified protein to an aldehyde modified protein

The hydrazide-modified protein prepared in EXAMPLE 15 was reacted with aldehyde-modified protein in an identical manner as described for the hydrazine-modified protein in EXAMPLE 12. Analysis by PAGE gel demonstrated similar efficiency as conjugation as observed in EXAMPLE 12.

Please amend the paragraph on page 51, lines 14-22, as follows:

EXAMPLE 21

Immobilization of Oxidized Horseradish peroxidase to Hydrazine-Modified Plates

Periodate-oxidized horseradish peroxidase (Pierce Chemical Co., Rockford, IL) is diluted to the desired concentration and added to 96-3456 well plates that had been modified to possess hydrazino groups as described above in EXAMPLE 9. The antibody solution is allowed to incubate for 2-18 hours followed by removal of the solution and washing with 0.5% [t]Tween® solution (twice) and buffer (twice).

Please amend the paragraph beginning on page 53, line 18, through page 54, line 8, as follows:

EXAMPLE 26

Preparation of a[n] bifunctional aromatic oxyammonium hydrochloride, -O-oxyammoniumnicotinate hydrochloride (SONH)

As shown in FIGURE 9, to a solution of 6-chloronicotinic acid (1 equivalent) in 80% aqueous ethanol is added hydroxylamine (500 equivalents) and the solution is refluxed for 16 hours. The reaction mixture is concentrated to dryness and dissolved in water. The solution cooled in an ice bath and acidified with concentrated hydrochloric acid until a precipitate forms, pH approximately 5.0. The solids are isolated, redissolved in water and the pH of the solution raised to 7.5 with base. Dioxane (1 volume) is added to the solution followed by the dropwise addition of di-t-butyl dicarbonate (5 equivalents; Aldrich Chemical Co.). The reaction mixture is stirred at room temperature for 4 hours and the dioxane removed on the rotavap. The residue is chromatographed on silica to isolate the desired BOC acid. The acid (1 equiv) is dissolved in DMF and treated with NHS (1 equiv) followed by the dropwise addition of DCC (1 equiv) in DMF. The reaction mixture is stirred at room temperature for 4 hours and the solids removed by filtration and the filtrate concentrated to dryness and resuspended in ethyl acetate. Further precipitate is removed by filtration and the filtrate concentrated to dryness and the desired BOC succinimidyl ester is isolated by silica gel chromatography.

Please amend the paragraphs beginning on page 54, line 18, through page 55, line 30, as follows:

EXAMPLE 27

Preparation of an oxyamino-modified protein

A 5 mg/mL solution of ovalbumin in PBS (100 mM phosphate, 150 mM NaCl, pH 7.4) and 2 mM EDTA (200 μ L; 1 mg protein) is prepared. A solution of succinimidyl aminooxyacetate hydrochloride (SAAH) (2 mg) in DMF (50 μ L) is prepared. To the protein solution is added the SAAH/DMF solution (15 eq.). The reaction mixture is incubated at room temperature for 4 hours. The modified protein is isolated and buffer exchanged by placing the reaction mixture in a 30K ultra-free centrifugation device and washing three times with 0.1 M MES, 0.9% NaCl, pH 4.7 (3 X 400 μ L). The purified protein is quantified for protein concentration (BCA assay; Pierce Chemical Co., Rockford, IL).

EXAMPLE 28

Preparation of a hydrazinonicotinamide modified polymer

A solution of poly-L-lysine (10 mg; Sigma Chemicals, St. Louis, MO; cat. #P-7890) was dissolved in conjugation buffer, 0.1 M phosphate, 0.15 M NaCl, pH 7.4 (1 mL). A solution of succinimidyl 6-hydrazinonicotinate acetone hydrazone (SANH; 1.3 mg) was dissolved in DMSO (13 μ L). To two poly-L-lysine aliquots (200 μ L) were added the SANH/DMSO solution (2.85 μ L (10 equivalents) and 5.7 μ L (20 equivalents)). The reaction mixtures were vortexed and incubated at room temperature for 2 h. The modified polymer was isolated by gel filtration on a NAP-25 column (Phar[m]acia) pre-equilibrated with 0.1 M MES, 0.9% NaCl, pH 4.7 buffer. Fractions (1 mL) were collected and analyzed by UV (A260). Fractions containing UV active product were combined to yield the desired product. The product was analyzed colorimetrically for hydrazine content by dissolving an aliquot (2 μ L) in a 0.5 mM solution of p-nitrobenzaldehyde (98 μ L) and incubating at 37 °C for 1 h followed by taking A390 readings (extinction coefficient 22000). The HyNic:poly-L-lysine polymer was used directly in the conjugation step. The amine/hydrazine content was

determined using the TNBSA assay (trinitrobenzenesulfonic acid; Pierce Chemical, Inc., Rockville, IL).

EXAMPLE 29

Preparation of an aliphatic hydrazine modified protein and conjugation to periodate oxidized bacterial polysaccharide

A solution of a bacterial polysaccharide that possesses unsaturation in its lipids (from ATCC; 10 mg/mL) in water is treated with 10 mM sodium periodate (1/10 volume to make the solution 1 mM in periodate) and incubated at room temperature for 30 min. The reaction mixture is passed through a sephadex G-25 column pre-equilibrated with water to remove small molecule impurities. The polysaccharide combining fractions are combined and concentrated to 5 mg/mL.

IN THE CLAIMS:

Please amend claims 12, 23, 24, 35, 37, and 49-53 as follows:


12. (Amended) The compound of claim 10, wherein R is a divalent aliphatic group.

23. (Amended) The compound of claim 1, wherein B is an amino reactive moiety selected from succinimidyl ester, hydroxybenzotriazolyl ester, or pentafluorophenol ester.

24. (Amended) The compound of claim 1, wherein B is a thiol reactive moiety selected from maleimido, α -bromoacetyl, α -bromoacetamido or pyridyldisulfide.

35. (Amended) A method of crosslinking a natural or synthetic biological molecule, comprising:

(i) preparing a conjugate of formula Va:



or a derivative thereof, wherein:

A is NH(C=O), NH(C=S), NH(C=NH), NHNH(C=O), NHNH(C=S), NHNH(C=NH) or a direct bond;

B is a natural or synthetic biological molecule;

D is a carbon or nitrogen atom;

E is a carbon or nitrogen atom;


R¹ is hydrogen or a saturated straight chain of 1 to 12 carbon atoms; and


R² is hydrogen or a saturated straight chain of 1 to 12 carbon atoms; and

(ii) applying the conjugate to a surface wherein the surface has at least one [amino or one thiol reactive] carbonyl moiety for a time and under conditions such that the hydrazine moiety of the conjugate reacts with the [amino or thiol reactive] carbonyl moiety of the surface forming a hydrazone bond to the surface.

37. (Amended) A method of crosslinking a natural or synthetic biological molecule, comprising:

(i) preparing a conjugate of formula IVa:





A is $\text{NH}(\text{C}=\text{O})$, $\text{NH}(\text{C}=\text{S})$, $\text{NH}(\text{C}=\text{NH})$, $\text{NHNH}(\text{C}=\text{O})$, $\text{NHNH}(\text{C}=\text{S})$, $\text{NHNH}(\text{C}=\text{NH})$ or a direct bond;

D is a carbon or nitrogen atom;

X is a negative counter ion, oxygen, sulfur or -NH; and

(ii) mixing the conjugate to a natural or synthetic biological molecule, wherein the molecule has at least one carbonyl moiety, for a time and under conditions such that the hydrazine moiety of the conjugate reacts with the carbonyl moiety of the molecule forming a hydrazone bond to the molecule.

49. (Amended) The compound of claim 5, wherein B is an amino reactive moiety selected from succ[in]inimidyl ester, hydroxybenzotriazolyl ester, or pentafluorophenol ester.

50. (Amended) The compound of claim 8, wherein B is an amino reactive moiety selected from succ[in]inimidyl ester, hydroxybenzotriazolyl ester, or pentafluorophenol ester.

51. (Amended) The compound of claim 10, wherein B is an amino reactive moiety selected from succ[in]inimidyl ester, hydroxybenzotriazolyl ester, or pentafluorophenol ester.

52. (Amended) The compound of claim 5, wherein B is a thiol reactive moiety selected from maleimido, α -bromoacetyl, α -bromoacetamido or pyridyldisulfide.

U.S.S.N 09/815,978
Schwartz
PRELIMINARY AMENDMENT ATTACHMENT

53. (Amended) The compound of claim 10, wherein B is a thiol reactive moiety selected from maleimido, α -bromoacetyl, α -bromoacetamido or pyridyldisulfide.

* * *

Respectfully submitted,
HELLER EHRMAN WHITE & McAULIFFE LLP

By:


Stephanie L. Seidman
Registration No. 33, 779

Attorney Docket No. 37154-753
Address all correspondence to:
Stephanie L. Seidman, Esq.
HELLER EHRMAN WHITE & McAULIFFE
4350 La Jolla Village Drive, 6th Floor
San Diego, California 92122-1246
Telephone: 858 450-8400
Facsimile: 858 587-5360
email:sseidman@HEWM.com